

Anti-Dibromo-Tyrosine (DiBrY) Monoclonal Antibody

- Code:** MBY-020P
- Content:** 20 μ g of IgG / vial. Lyophilized powder.
- Clone #:** 3A5
- Immunogen:** Brominated keyhole limpet hemocyanin.
- Subclass:** Mouse IgG₁(κ), Protein-A purified.
- Application:** Immunohistochemistry (Recommended concentration: 10 μ g/mL IgG) [ref.1] and western blotting.
- Reconstitution:** Reconstitute with 200 μ L of distilled water.
- Buffer Concentration:** 100 μ g/mL of IgG in 10mM Phosphate buffered saline (pH7.4).
- Specificity:** Specific for di-halogenated tyrosine.
Recognize dibromotyrosine (DiBrY), N-acetylated dichloro tyrosine (NAcDiClY), diiodo tyrosine (DiIY), N-bromosuccinimide-treated BSA, HOCl-treated BSA and dibromo-tyrosone conjugated BSA.
- Cross reactivity was checked for dityrosine-conjugated BSA, peroxydinitrite-treated BSA, tyrosine (Y), chlorotyrosine(ClY), 3-3'-dityrosine (DiY), 3-aminotyrosine (NH₂Y), 3-nitrotyrosine (NO₂Y), phenylalanine (Phe), o-phosphotyrosine, 3,4-dihydroxyphenylacetic acid (DOPA) and 2-hydroxy-4-phenylacetic acid. Bromotyrosine (BrY) is weakly recognized.
- Storage:** Store at less than -20°C.
After reconstitution, store as frozen aliquots and avoid repeated freeze & thaw.
- Stability:** 3 years at -20°C.
- References:** 1) Y Kato, Y Kawai, H Morinaga, H Kondo, N Dozaki, N Kitamoto and T Osawa.: Immunogenicity of a brominated protein and successive establishment of a monoclonal antibody to dihalogenated tyrosine. Free Rad Biol Med, 38, p24-31 (2005)

For research use only, not for diagnostic use.



JAPAN INSTITUTE FOR THE CONTROL OF AGING (JaICA)

710-1 Haruoka, Fukuroi, Shizuoka, 437-0122 Japan

TEL : +81-538-49-0125

FAX : +81-538-49-1267

WEB : <http://www.jaica.com>

E-mail : biotech@jaica.com